

L. Podda, F. Meloni, G. Fenu, G. Iiriti & G. Bacchetta

The vascular flora of the Marine Protected Area of “Capo Carbonara” (SE-Sardinia)

Abstract

Podda, L., Meloni, F., Fenu, G., Iiriti, G. & Bacchetta, G.: The vascular flora of the Marine Protected Area of “Capo Carbonara” (SE-Sardinia). — *Fl. Medit.* 31 (Special Issue): 415-449. 2021. — ISSN: 1120-4052 printed, 2240-4538 online.

This study aims at presenting an updated inventory of the vascular flora of the Capo Carbonara Marine Protected Area, including the Sites of Community Importance “Isola dei Cavoli, Serpentara, Punta Molentis e Campulongu” and partially of “Costa di Cagliari”, in Sardinia (Italy). A total of 653 taxa are reported, 84% of which are native (5% endemics) and 16% non-native (10% neophytes, 6% archaeophytes). The native component includes 406 species, 135 subspecies, two variety and one hybrid, belonging to 75 families and 313 genera. The non-native taxa are 94 species, 13 subspecies, one variety and one hybrid, belonging to 40 families and 81 genera.

Life-form analysis of native species revealed a prevalence of therophytes (49%) followed by hemicryptophytes (21%), geophytes (14%) and chamaephytes (7%). Therophytes (33%) followed by phanerophytes (32%) are prevalent among the non-native taxa. Biogeographically, among native species, the Mediterranean element is largely prevailing (81%), mainly consisting of Circum-Mediterranean (39%) and Euro-Mediterranean (27%) taxa, while the American element (35%) prevails over the Mediterranean species (29%) among the non-native taxa. Among endemics, the Sardo-Corsican taxa are dominant (55%), followed by Sardo-Corsican-Tuscan Archipelago (14%) and Sardinian (10%) taxa.

Key words: Alien species, biodiversity, conservation, endemics, marine protected areas, Mediterranean Basin, Sardinia, vascular flora.

Introduction

The territory of the Marine Protected Area of Capo Carbonara (hereafter MPACC) is known as one of the most important coastal areas in Sardinia. Since 1998, the year of its legal establishment, it is also the only marine protected area in southern Sardinia. The institution of the MPACC, together with the previous establishment of the Sites of Community Importance (SCI) and Special Protection Areas (SPA) of the Natura 2000 Network, increased the protection level of this territory, which is very fragile due to touristic pressure, and have triggered several biodiversity conservation projects (AA.VV. 2014; Acunto & al. 2017; Pinna & al. 2017).

The territory shows a remarkable floristic diversity due to the richness of habitats and has attracted several botanists since the early 1800s. The first reports about the vascular flora of this area, from Capo Carbonara and Porto Giunco, are due to Moris (1827, 1837-1859). Subsequently, between 1894 and 1916, Martelli carried out some herborizations in different sites of south-eastern Sardinia, including the Isola dei Cavoli. The area of Capo Carbonara was frequently visited by botanists because it was accessible by a coastal road which, although defined at that time as very dangerous, allowed to reach this area (Bocchieri & Iiriti 2007a).

Some herborizations along the coast of Villasimius were carried out by Martinoli between 1940 and 1951 (Bocchieri & Iiriti 2007a). More recent investigations have focused on the Cavoli (Corsi 1963; Mossa & Tamponi 1978; Mossa & Fogu 1987) and Serpentara (Bocchieri 1988, 1989; Biondi & al. 1993) islands, Capo Carbonara, and the surrounding areas (Camarda & Ballero 1981; Mossa & al. 2000). Further floristic information for the study area are reported in studies addressing areas of south-eastern Sardinia or carried out at a regional scale (Chiappini & Diana 1978; Ballero 1982; Arrigoni & Bocchieri 1995; Mayer 1995; Bocchieri 2001; Palmese & al. 2001; Mossa & al. 2003; Bacchetta & al. 2005; Arrigoni 2006-2015; Iiriti 2006; Bocchieri & Iiriti 2007a, 2007b, 2008).

To date, however, there is no updated checklist of the vascular flora of this area (Bacchetta & al. 2006, 2007; Bacchetta & al. 2015; Pinna & al. 2017). In the framework of the project “Conservation of Plant Biodiversity in the Marine Protected Area of Capo Carbonara”, funded by the Ministry of the Environment and Territory and Sea Protection (MATM) (Bacchetta & al. 2006, 2007), the project LIFE07 NAT/IT/000519 PROVIDUNE (Durán & al. 2016; Pinna & al. 2015, 2017) and LIFE13 NAT/IT/000433 RES MARIS project (Acunto & al. 2017; Bacchetta & al. 2018), in the years from 2014 to 2018, a study was carried out on the vascular flora of the coastal territory of the municipality of Villasimius, which includes the SCI “Isola dei Cavoli, Serpentara, Punta Molentis and Campulongu”, coinciding with MPACC.

This study brings an overview of the vascular flora in the MPACC and contributes to update the information on the native and exotic flora, thus providing a basis for conservation actions.

Study area

The study area includes coastal and insular areas of the MPACC in south-eastern Sardinia (Fig. 1). The MPACC, 8.598 hectares in surface, was founded by a Ministerial Decree on 15/09/98 (modified on 3/08/99) with the participation of the Sardinian Region and Villasimius municipality. It encloses the marine area in front of the Cavoli and Serpentara islands and the coast belonging to Villasimius. The coast is 41 km long and includes several beaches as Porto Giunco, Simius, Campu Longu and Campus, where rocky promontories alternate with small bays such as Porto Sa Ruxi or Punta Molentis.

The sea areas are characterized by a continuity of granite forms with emerging spots, while wide sandy expanses connect the islands interrupted by granitic shoals.

The lithologies are essentially referred to the Ercinic batholith of Sarrabus emerged at the end of the Ercinic orogenesis in the Middle Carboniferous (Carmignani & al. 1982). Rocks mainly




Fig. 1. Geographic position of the Marine Protected Area of Capo Carbonara (MPACC).

derive from granites and granodiorites subjected, during the alpine orogenesis, to tectonic movements that deformed then according to a NW-SE prevalent direction. The veins present the same orientation. They are mainly of a basic nature, dominated by lamprophyric and spessartitic lithologies which are important in shaping the coastal and submarine landscape, the coastline being formed by promontories with the same orientation as those of Capo Carbonara (Orrù & al. 1994). The oldest organogenic deposits are formed by fossiliferous beaches conglomerates known as Tyrrhenian Platform, which appear in various points of Capo Carbonara promontory. One of the most important outcrops is near Cava Usai, in the locality of Portu su Forru, which has a total extension of about 400 m. This formation is surmounted by sandstones of aeolic origin (paleodunes), a reddish paleosoil (palexeralf), and, in the submerged environment, by the sedimentation and cementation of beach-rocks. In the surrounding landscape, ancient shapes of smoothing such as “inselberg” and “tor” ornament the ridges and the plateaux at an elevation ranging from 400 to 500 m (Orrù & al. 1994).

Climate is typically Mediterranean; thermometric data (37 years of observations) from the climatic station of Capo Carbonara, show a strong seasonal trend in temperature, typical of the coast of Sardinia. Annual mean temperature is 21.8°C, mean maximum temperature of the hottest month 30.9°C (August) and mean minimum temperature of the coldest month 9.5°C (February). As for precipitation, the climatic station of Capo Carbonara recorded an annual mean of 238.7 mm, with higher values in autumn; October, with 35 mm on average, is the wettest month (Table 1).

Table 1. Mean monthly and annual temperature (°C) and rainfall (mm), Capo Carbonara station (Italian Air Force National Meteorological Service). Temperature observations: 1960-2007 (37 years); rainfall observations: 1954-2008 (54 years).

	Jan	Feb.	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Mean Temp	11.6	11.6	12.7	14.4	18.1	22.2	25.3	26.0	23.3	19.8	15.9	12.8	17.8
Max Temp	14.7	14.9	16.6	18.2	22.2	26.7	30.2	30.9	27.6	24.0	19.2	15.7	21.8
Min Temp	9.9	9.5	11.0	12.3	15.8	19.8	22.8	23.8	21.0	18.0	14.1	11.0	15.7
Rainfall	22.6	24.3	23.3	20.0	13.0	3.4	2.5	7.3	27.7	35.0	33.0	26.7	238.7

From a bioclimatic point of view, the study area has a Mediterranean Pluviseasonal Oceanic (MPO) bioclimate, a Thermo-Mediterranean (TME) thermotype and an ombrotype ranging from upper semiarid to lower dry (Bacchetta & al. 2009).

The superficial movable bottom is mainly colonized by beds of *Posidonia oceanica* Delile, being a part of the priority habitat 1120 “*Posidonia* beds (*Posidonion oceanicae*)” (Acunto & al. 2017).

The dune ecosystems are well preserved, hosting in different places the complete psamphilous geosigmatata from the annual herbaceous vegetation to the micro-forest dominated by junipers of the priority habitat 2250 “Coastal dunes with *Juniperus* spp.” and pine of the priority habitat 2270 “Wooded dunes with *Pinus pinea* and/or *Pinus pinaster*” (Acunto & al. 2017). The populations of *Crucianella maritima* L. are of a great value. Even if fragmented, they occupy the back side of the dunes and belong to the habitat 2210 “*Crucianellion maritimae* fixed beach dunes”, as well as the formations of *Achillea maritima* (L.) Ehrend. & Y.P.Guo that can be found only in Cava Usai, Punta Molentis and a few other places along the south-eastern coast of Sardinia. The rocky surfaces and cliffs in front of the sea are occupied by the association *Crithmo maritimi-Limonietum retiramei corr.* (Mossa & Tamponi 1978) attributable to the habitat 1240 “Cliffs with the Mediterranean coastal vegetation with *Limonium* spp. endemic”. The Notteri pond occupies the back side of the dune area of the Porto Giunco beach. It occupies a surface of about 34 ha and hosts an interesting halophyllous vegetation referred to the priority habitat 1510 “Mediterranean salt steppes (*Limonietalia*)”, while the muddy bottom is colonized by the association *Chaetomorpha-Ruppium* Br.-Bl. 1952 being a part of the habitat 1150 “Coastal lagoons” (Mossa & al. 2000).

In the framework of the Italo-Tyrrhenian biogeographic superprovince (Ladero-Alvarez 1987; Bacchetta & al. 2013) and the Sardinian-Corsican biogeographic province (Bacchetta & al. 2012), based on the vascular endemic flora, the study area falls into the Campidanese-Turritano sector and Sarrabense subsector (Fenu & al. 2014).

Material and Methods

In order to perform the analysis of the vascular flora, floristic data retrieved from literature were implemented by a four years' work that included field investigation during the MATTM, LIFE PROVIDUNE and LIFE RES MARIS projects, and the revision of herbarium and bibliographic material.

The taxonomic treatment of plant species and subspecies follow the last checklists of the native and alien floras of Italy (Bartolucci & al. 2018; Galasso & al. 2018). Plant families have been validated according to the Angiosperm Phylogeny Group IV (APG 2016).

Life-forms were directly checked in the field and expressed according to the Raunkiaer's classification system (Raunkiaer 1934), using the abbreviations reported in Pignatti (1982). For the biogeographical types we used Pignatti's classification (Pignatti 1982), as modified by Brullo & al. (1996) for Mediterranean chorotypes. For endemic taxa, the categories proposed by Arrigoni & Di Tommaso (1991) were adopted, as modified by Bacchetta & Pontecorvo (2005).

The geographic origin of alien plants was based on Podda & al. (2012) and Puddu & al. (2016). The status of alien species was determined and ordered according to Richardson & al. (2000), Pyšek & al. (2004), and Richardson & al. (2011). Archaeophyte and neophyte taxa were differentiated depending on their introduction before or after the years 1492/1500, respectively.

Floristic checklist

Db = dubious reporting

Alien taxa: Neo = Neophytes; Arch = Archeophytes; Arch D = cryptogenic; Status alien taxa: Inv = invasive; Nat = naturalised; Cas = casual.

PTERIDOPHYTA

SELAGINELLACEAE

Selaginella denticulata (L.) Spring – Ch rept – Circum-Medit.

EQUISETACEAE

Equisetum ramosissimum Desf. – G rhiz – Circumbor.

DENNSTAEDTIACEAE

Pteridium aquilinum (L.) Kuhn subsp. *aquilinum* – G rhiz – Cosmop.

ASPLENIACEAE

Asplenium ceterach L. subsp. *ceterach* – H ros – Eurasiat.

Asplenium obovatum subsp. *billotii* (F.W.Schultz) O.Bolòs, Vigo, Masalles & Ninot – H ros – W-Medit.

Asplenium obovatum Viv. subsp. *obovatum* – H ros – Circum-Medit.

PTERIDACEAE

Anogramma leptophylla (L.) Link – T caesp – Subtrop.

Oeosporangium pteridioides (Reichard) Fraser-Jenk. & Pariyar – H ros – W-Medit.

POLYPODIACEAE

Polypodium cambricum L. – H ros – Euro-Medit.

PINOPHYTA

CUPRESSACEAE

Cupressus sempervirens L. – P scap – E-Medit. (Arch – Cas)

Juniperus macrocarpa Sm. – P caesp – Circum-Medit.

Juniperus turbinata Guss. – P caesp – Euro-Medit.

PINACEAE

Pinus halepensis Mill. subsp. *halepensis* – P scap – Circum-Medit.

Pinus pinea L. – P scap – Euri-Medit. (Arch – Nat)

MAGNOLIOPHYTA

Magnolides

ARISTOLOCHIACEAE

Aristolochia rotunda subsp. *insularis* (E. Nardi & Arrigoni) Gamisans – G rhiz – Endem.
SA-CO

Aristolochia tyrrhena E.Nardi & Arrigoni – G rad – Endem. SA

Monocotyledones

ARACEAE

Ambrosina bassii L. – G rhiz – SW-Medit.

Arisarum vulgare O.Targ.-Tozz. subsp. *vulgare* – G rhiz – Circum-Medit.

Arum pictum L.f. subsp. *pictum* – G rhiz – Endem. SA-CO-AT

Helicodiceros muscivorus (L.f.) Engl. – G rhiz – Endem. SA-CO-BL

Lemna minor L. – I nat – Boreo-Trop.

CYMODOCEACEAE

Cymodocea nodosa (Ucria) Asch. – I rad – Medit.-Atl.

POTAMOGETONACEAE

Posidonia oceanica (L.) Delile – I rad – Circum-Medit.

Ruppia maritima L. – I rad – Boreo-Trop.

AMARYRILLIDACEAE

Acis autumnalis (L.) Sweet – G bulb – Circum-Medit.

Allium cepa L. – G bulb – Asia. (Arch – Cas)

Allium commutatum Guss. – G bulb – Circum-Medit.

Allium polyanthum Schult. & Schult.f. – G bulb – Euro-Medit.

Allium roseum L. subsp. *roseum* – G bulb – Circum-Medit.

- Allium sativum* L. – G bulb – Europ. (Arch – Cas)
Allium subhirsutum L. subsp. *subhirsutum* – G bulb – Circum-Medit.
Allium triquetrum L. – G bulb – Circum-Medit.
Narcissus miniatus Donn.-Morg., Koop. & Zonn. – G bulb – Circum -Medit.
Narcissus supramontanus subsp. *cunicularium* Arrigoni – G bulb – Endem. SA-CO
Pancratium maritimum L. – G bulb – Circum-Medit.

ASPARAGACEAE

- Agave americana* L. – P caesp – Neotrop. (Neo – Inv)
Agave attenuata Salm-Dyck subsp. *attenuata* – P caesp – Neotrop. (Neo – Cas)
Agave fourcroydes Lem. – P caesp – Neotrop. (Neo – Inv)
Agave ingens Brg. var. *picta* – P caesp – Neotrop. (Neo – Inv)
Agave salmiana Otto ex Salm-Dyck subsp. *ferox* (K.Koch) Hochstätter – P caesp – Neotrop. (Neo – Inv)
Asparagus acutifolius L. – G rhiz – Circum-Medit.
Asparagus albus L. – Ch frut – W-Medit.
Asparagus aphyllus L. – Ch frut – S-Medit.
Asparagus horridus L. – NP – S-Medit.
Charybdis pancration (Steinh.) Speta – G bulb – Circum-Medit.
Charybdis undulata (Desf.) Speta – G bulb – S-Medit.
Muscari comosum (L.) Mill. – G bulb – Euro-Medit.
Prospero corsicum (Boullu) J.-M.Tison – G bulb – Endem. SA-CO
Prospero obtusifolium subsp. *intermedium* (Guss.) Soldano & F.Conti – G bulb – SW-Medit.
Ruscus aculeatus L. – Ch frut – Euro-Medit.
Ruscus hypoglossum L. – G rhiz – Euro-Medit. (Neo – Cas)
Urginea fugax (Moris) Steinh. – G bulb – SW-Medit.
Yucca aloifolia L. – P caesp – Neotrop. (Neo – Cas)
Yucca gloriosa L. – P caesp – Neotrop. (Neo – Cas)

IRIDACEAE

- Chamaeiris foetidissima* (L.) Medik. – G rhiz – Euro-Medit.
Gladiolus byzantinus Mill. – G bulb – Circum-Medit. (Arch D – Nat)
Iris germanica L. – G rhiz – Europ. (Arch – Nat)
Limniris pseudacorus (L.) Fuss – G rhiz – Eurasiat.
Moraea sisyrinchium (L.) Ker Gawl. – G bulb – Circum-Medit.
Romulea columnae Sebast. & Mauri – G bulb – Circum-Medit.
Romulea ligustica Parl. – G bulb – SW-Medit.
Romulea ramiflora Ten. subsp. *ramiflora* – G bulb – Circum-Medit.
Romulea requienii Parl. – G bulb – Endem. SA-CO
Romulea rollii Parl. – G bulb – W-Medit.
Xiphion xiphium (L.) M.B. Crespo, Mart.Azorín & Mavrodie – G bulb – W-Medit. (Arch Db – Nat)

ORCHIDACEAE

Anacamptis collina (Banks & Sol. ex Russell) R.M.Bateman, Pridgeon & M.W.Chase – G bulb – Circum-Medit.

Anacamptis laxiflora (Lam.) R.M. Bateman, Pridgeon & M.W. Chase – G bulb – Circum-Medit.

Anacamptis longicornu (Poir.) R.M. Bateman, Pridgeon & M.W. Chase – G bulb – W-Medit.

Anacamptis papilionacea (L.) R.M. Bateman, Pridgeon & M.W. Chase – G bulb – W-Medit.

Ophrys apifera Huds. – G bulb – Euro-Medit.

Ophrys bombyliflora Link – G bulb – Circum-Medit.

Ophrys conradiæ Melki & Deschâtres – G bulb – Endem. SA-CO

Ophrys exaltata subsp. *morisii* (Martelli) Del Prete – G bulb – Endem. SA-CO

Ophrys incubacea Bianca – G bulb – W-Medit.

Ophrys speculum Link – G bulb – Circum-Medit.

Ophrys tenthredinifera Willd. subsp. *neglecta* (Parl.) E.G.Camus – G bulb – N-Medit.

Orchis × *bornemanniae* Asch. – G bulb – W-Medit.

Serapias lingua L. – G bulb – Medit.-Atl.

Serapias parviflora Parl. – G bulb – Medit.-Atl.

Spiranthes spiralis (L.) Chevall. – G rhiz – Europ.-Caucas.

ASPHODELACEAE

Aloe arborescens Mill. – NP – S-Africa (Neo – Cas)

Aloe × *caesia* Salm-Dyck – NP – Africa (Neo – Cas)

Asphodelus fistulosus L. – H bien – Circum-Medit.

Asphodelus ramosus L. subsp. *ramosus* – G rhiz – Circum-Medit.

DIOSCOREACEAE

Dioscorea communis (L.) Caddick & Wilkin. – G rad – Euro-Medit.

SMILACACEAE

Smilax aspera L. – P lian – Circum-Medit.

ARECACEAE

Phoenix canariensis H.Wildpret – P scap – Macarones. (Neo – Cas)

Washingtonia filifera (Linden ex André) H.Wendl. ex de Bary – P scap – Neotrop. (Neo – Cas)

CYPERACEAE

Carex distachya Desf. – H caesp – Circum-Medit.

Carex divisa Huds. – G rhiz – Medit.-Atl.

Carex flacca subsp. *erythrostachys* (Hoppe) Holub – G rhiz – Circum-Medit.

Carex halleriana Asso – H caesp – Euro-Medit.

Cyperus capitatus Vand. – G rhiz – Circum-Medit.

Cyperus rotundus L. – G rhiz – Medit-Trop. (Arch D – Nat)

Schoenus nigricans L. – H caesp – Subcosmop.

Scirpoides holoschoenus (L.) Soják – G rhiz – Euro-Medit.

*JUNCACEAE**Juncus acutus* L. subsp. *acutus* – H caesp – Euro-Medit.*Juncus bufonius* L. – T caesp – Cosmop.*Juncus maritimus* Lam. – G rhiz – Euro-Medit.*Juncus subulatus* Forssk. – G rhiz – S-Medit.*JUNCAGINACEAE**Triglochin barrelieri* Loisel. – G bulb – Circum-Medit.*POACEAE**Aira elegantissima* Schur subsp. *elegantissima* – T scap – Euro-Medit.*Ampelodesmos mauritanicus* (Poir.) T. Durand & Schinz – H caesp – SW-Medit.*Anisantha diandra* (Roth) Tutin ex Tzvelev – T scap – Euro-Medit.*Anisantha madritensis* (L.) Nevski subsp. *madritensis* – T scap – Medit.-Atl.*Anisantha rubens* (L.) Nevski – T scap – S-Medit.-Turan.*Anisantha sterilis* (L.) Nevski – T scap – Paleotemp.*Arundo donax* L. – G rhiz – Irano-Turan. (Arch – Inv)*Avena barbata* Pott ex Link – T scap – Euro-Medit.-Turan.*Avena fatua* L. – T scap – Irano-Turan. (Arch – Nat).*Avena sativa* L. subsp. *sativa* – T scap – Irano-Turan. (Arch – Cas)*Avena sterilis* L. – T scap – Circum-Medit. (Arch – Nat)*Brachypodium distachyon* (L.) P. Beauv. – T scap – Medit.-Turan.*Brachypodium retusum* (Pers.) P. Beauv. – H caesp – W-Medit.*Brachypodium sylvaticum* (Huds.) P. Beauv. subsp. *sylvaticum* – H caesp – Paleotemp.*Briza maxima* L. – T scap – Subtrop.*Briza minor* L. – T scap – Subcosmop.*Bromus hordeaceus* L. subsp. *hordeaceus* – T scap – Subcosmop.*Calamagrostis arenaria* subsp. *arundinacea* (Husn.) Banfi, Galasso & Bartolucci – G rhiz
– Circum-Medit.*Castellia tuberculosa* (Moris) Bor – T scap – Medit.-Irano-Turan. (Db)*Catapodium balearicum* (Willk.) H. Scholz – T scap – Medit.-Atl.*Catapodium rigidum* (L.) C.E. Hubb. subsp. *rigidum* – T scap – Euro-Medit.*Cenchrus longisetus* M.C. Johnst. – H caesp – S-Medit. (Neo – Nat)*Cortaderia selloana* (Schult. & Schult. f.) Asch. & Graebn. – H caesp – S-Amer. (Neo – Cas)*Cutandia divaricata* (Desf.) Barbey – T scap – Circum-Medit.*Cutandia maritima* (L.) Benth. ex Barbey – T scap – Circum-Medit.*Cynodon dactylon* (L.) Pers. – H rept – Subcosmop.*Cynosurus echinatus* L. – T scap – Euro-Medit.*Dactylis glomerata* subsp. *hispanica* (Roth) Nyman – H caesp – Circum-Medit.*Dasypyrum villosum* (L.) P. Candargy – T scap – Euro-Medit.-Turan.*Elymus repens* (L.) Gould subsp. *repens* – G rhiz – Paleotemp.*Festuca danthonii* Asch. & Graebn. subsp. *danthonii* – T scap – Euro-Medit.*Festuca fasciculata* Forssk. – T caesp – Circum-Medit.*Festuca geniculata* (L.) Lag. & Rodr. subsp. *geniculata* – T caesp – W-Medit.*Festuca ligustica* (All.) Bertol. – T caesp – Circum-Medit.

- Gastridium ventricosum* (Gouan) Schinz & Thell. – T scap – Medit.-Atl.
Holcus annuus subsp. *setiglumis* (Boiss. & Reut) M.Seq. & Castrov. – T scap – Circum-Medit.
Hordeum murinum subsp. *leporinum* (Link) Arcang. – T scap – Euro-Medit.
Hyparrhenia hirta (L.) Stapf subsp. *hirta* – H caesp – Paleotrop.
Imperata cylindrica (L.) Raeusch. – G rhiz – Cosmop.
Lagurus ovatus L. subsp. *ovatus* – T scap – Euro-Medit.
Lagurus ovatus L. subsp. *vestitus* (Messeri) Brullo – T scap – S-Medit.
Lamarckia aurea (L.) Moench – T scap – Medit.-Turan.
Lolium rigidum Gaudin – T scap – Paleo-Subtrop.
Lygeum spartum L. – H caesp – S-Medit.
Melica ciliata L. subsp. *ciliata* – H caesp – Euro-Medit.
Melica minuta L. – H caesp – Circum-Medit.
Oloptum thomasi (Duby) Banfi & Galasso – H caesp – Circum-Medit.
Parapholis incurva (L.) C.E.Hubb. subsp. *incurva* – T scap – Medit.-Atl.
Parapholis strigosa (Dumort.) C.E.Hubb. – T scap – Medit.-Atl.
Paspalum distichum L. – G rhiz – Neotrop. (Neo – Nat)
Phalaris brachystachys Link – T scap – Circum-Medit.
Phalaris coerulescens Desf. – H caesp – Medit.-Macarones.
Phalaris minor Retz. – T scap – Medit.-Irano-Turan.
Phleum arenarium subsp. *caesium* H.Scholz – T scap – Circum-Medit.
Phragmites australis (Cav.) Trin. ex Steud. – G rhiz – Subcosmop.
Piptatherum coerulescens (Desf.) P. Beauv. – H caesp – Circum-Medit.
Poa annua L. – T caesp – Cosmop.
Poa bulbosa L. subsp. *bulbosa* – G bulb – Paleotemp.
Polypogon monspeliensis (L.) Desf. – T scap – Subtrop.
Polypogon subspathaceus Req. – T scap – Circum-Medit.
Rostraria cristata (L.) Tzvelev – T scap – Medit.-Irano-Turan.
Rostraria pubescens (Lam.) Trin. – T scap – Circum-Medit.
Sporobolus pumilus (Roth) P.M.Peterson & Saarela – G rhiz – Anfiatl. (Neo – Nat)
Sporobolus virginicus (L.) Kunth – G rhiz – Circum-Medit.
Stipellula capensis (Thunb.) Röser & H.R.Hamasha – T scap – Circum-Medit.
Thinopyrum elongatum (Host) D.R.Dewey – H caesp – Euro-Medit.
Thinopyrum junceum (L.) Á.Löve – G rhiz – Circum-Medit.
Triticum neglectum (Req. ex Bertol.) Greuter – T scap – Medit.-Turan.
Triticum ventricosum (Tausch) Ces., Pass. & Gibelli – T scap – W-Medit. (Arch D – Nat)

TYPHACEAE

- Typha angustifolia* L. – G rhiz – Circumbor.

CANNACEAE

- Canna indica* L. – G rhiz – Neotrop. (Neo-Cas)

Eudicotyledones

PAPAVERACEAE

Fumaria capreolata L. subsp. **capreolata** – T scap – Euro-Medit.
Fumaria flabellata Gasp. – T scap – Circum-Medit.
Fumaria officinalis L. subsp. **officinalis** – T scap – Subcosmop.
Glaucium flavum Crantz – H scap – Circumbor.
Hypecoum procumbens L. subsp. **procumbens** – T scap – Paleotemp.
Papaver hybridum L. – T scap – Medit.-Turan.
Papaver rhoeas L. subsp. **rhoeas** – T scap – E-Medit. (Arch D – Nat)

RANUNCULACEAE

Anemone hortensis L. subsp. **hortensis** – G bulb – N-Medit.
Clematis cirrhosa L. – P lian – Medit.-Turan.
Clematis flammula L. – P lian – Euro-Medit.
Ficaria verna subsp. **fertilis** (Lawalrée ex Laegaard) Stace – G bulb – Euroasiat.
Nigella damascena L. – T scap – Euro-Medit.
Ranunculus bullatus L. – H ros – Circum-Medit.
Ranunculus paludosus Poir. – H scap – Medit.-Turan.

AMARANTHACEAE

Achyranthes sicula (L.) All. – Ch suffr – W-Medit.
Amaranthus albus L. – T scap – Neotrop. (Neo – Inv)
Amaranthus blitum L. subsp. **blitum** – T scap – Boreo-Trop.

CHENOPODIACEAE

Arthrocaulon macrostachyum (Moric.) Piirainen & G.Kadereit – Ch succ – Medit.-Irano-Turan. (Db)
Atriplex prostrata Boucher ex DC. – T scap – Circumbor.
Beta vulgaris L. subsp. **maritima** (L.) Arcang. – T scap – Euro-Medit.
Beta vulgaris L. subsp. **vulgaris** – H scap – Euro-Medit. (Arch – Cas)
Chenopodium album L. subsp. **album** – T scap – Subcosmop.
Chenopodium murale (L.) S.Fuentes, Uotila & Borsch – T scap – Subcosmop.
Halimione portulacoides (L.) Aellen – Ch frut – Circumbor.
Oxybasis glauca (L.) S. Fuentes, Uotila & Borsch – T scap – Subcosmop. (Db)
Salicornia fruticosa (L.) L. – Ch succ – Boreo-Trop.
Salsola squarrosa subsp. **controversa** (Tod. ex Lojac.) Mosyakin – T scap – Circumbor.
Soda inermis Fourr. – T scap – Paleotemp.
Suaeda vera J.F.Gmel. – Ch suffr – Subcosmop.

AIZOACEAE

Carpobrotus acinaciformis (L.) L.Bolus – Ch suffr – S-Africa (Neo – Inv)
Carpobrotus edulis (L.) N.E.Br. – Ch suffr – S-Africa (Neo – Inv)
Drosanthemum floribundum (Haw.) Schwantes – H scand – S-Africa (Neo – Nat)
Malephora crocea (Jacq.) Schwantes – Ch scap – S-Africa (Neo – Inv)
Mesembryanthemum cordifolium L.f. – H rept – S-Africa (Neo – Inv)
Mesembryanthemum cristallinum L. – T scap – S-Africa (Neo – Nat)
Mesembryanthemum nodiflorum L. – T scap – S-Africa (Neo – Nat)

CACTACEAE

- Austrocylindropuntia subulata* (Mühlenpf.) Backeb. – P succ – Neotrop. (Neo – Inv)
Nopalea dejecta Salm-Dick – P succ – Neotrop. (Neo – Cas)
Opuntia dillenii (Ker Gawl.) Haw. – P succ – Neotrop. (Neo – Nat)
Opuntia ficus-indica (L.) Mill. – P succ – Neotrop. (Neo – Inv)
Opuntia puberula Hort. Vindob. ex Pfeiff. – P succ – Neotrop. (Neo – Cas)
Opuntia monacantha Haw. – P succ – Neotrop. (Neo – Nat)

CARYOPHYLLACEAE

- Arenaria leptoclados* (Rchb.) Guss. subsp. *leptoclados* – T scap – Paleotemp.
Cerastium glomeratum Thuill. – T scap – Euro-Medit.
Cerastium pumilum Curtis – T scap – Euro-Medit.
Corrigiola telephiifolia Pourr. – H ros – W-Medit.
Eudianthe coeli-rosa (L.) Fenzl ex Endl. – T scap – W-Medit.
Paronychia argentea Lam. – H caesp – Circum-Medit.
Petrorhagia dubia (Raf.) G.López & Romo – T scap – S-Medit.
Petrorhagia nanteuillii (Burnat) P.W.Ball & Heywood – T scap – W-Medit.
Petrorhagia prolifera (L.) P.W.Ball & Heywood – T scap – Euro-Medit.
Polycarpon tetraphyllum subsp. *alsinifolium* (Biv.) Ball – T scap – S-Medit.
Polycarpon tetraphyllum subsp. *diphyllum* (Cav.) O. Bolòs & Font-Quer – T scap – Euro-Medit.
Rhodalsine geniculata (Poiret) F.N.Williams – Ch suffr – W-Medit.
Sagina apetala Ard. subsp. *apetala* – T scap – Euro-Medit.
Sagina maritima Don – T scap – Medit.-Atl.
Silene bellidifolia Jacq. – T scap – S-Medit.
Silene colorata Poir. – T scap – Circum-Medit.
Silene gallica L. – T scap – Euro-Medit.
Silene niceensis All. – T scap – Circum-Medit.
Silene succulenta subsp. *corsica* (DC.) Nyman – H caesp – Endem. SA-CO
Silene valsecchiae Bocchieri – T scap – Endem. SA
Silene vulgaris subsp. *tenoreana* (Colla) Soldano & F. Conti – H caesp – Circum-Medit.
Silene vulgaris (Moench) Garcke subsp. *vulgaris* – H caesp – Circum-Medit.
Spergularia marina (L.) Besser – T scap – Paleotemp.
Spergularia rubra (L.) J.Presl & C. Presl – Ch suffr – Subcosmop.
Stellaria media (L.) Vill. subsp. *media* – T rept – Cosmop.
Stellaria neglecta Weihe subsp. *neglecta* – T scap – Paleotemp.
Stellaria pallida (Dumort.) Crép. – T scap – Euro-Medit.
Velezia rigida L. – T scap – Medit.-Turan.

FRANKENIACEAE

- Frankenia hirsuta* L. – Ch suffr – Subcosmop.
Frankenia laevis L. subsp. *laevis* – Ch suffr – Subcosmop.

NYCTAGINACEAE

- Mirabilis jalapa* L. – T caesp – Neotrop. (Neo – Inv)

PHYTOLACCACEAE

Phytolacca americana L. – G rhiz – Neotrop. (Neo – Nat)

PLUMBAGINACEAE

Limonium dubium (Andrews ex Guss.) Litard. – Ch suffr – Endem. SA-CO-SI

Limonium narbonense Mill. – H ros – Circum-Medit.

Limonium retirameum Greuter & Burdet subsp. ***retirameum*** – Ch suffr – Endem. SA

Plumbago europaea L. – Ch frut – Circum-Medit.

POLYGONACEAE

Polygonum maritimum L. – Ch rept – Subcosmop.

Rumex bucephalophorus L. subsp. ***bucephalophorus*** – T scap – Medit.-Macarones.

Rumex crispus L. – H scap – Paleotemp.

Rumex pulcher L. subsp. ***pulcher*** – H scap – Euro-Medit.

Rumex spinosus L. – T scap – Circum-Medit.

PORTULACACEAE

Portulaca sardoa Danin, Bagella & Marrosu – T scap – Endem. SA-CO

TAMARICACEAE

Tamarix africana Poir. – P scap – W-Medit.

Tamarix gallica L. – P caesp – W-Medit.

SANTALACEAE

Osyris alba L. – NP – Euro-Medit.

Thesium humile Vahl – T scap – Medit.-Atl.

CRASSULACEAE

Aeonium haworthii (Webb & Berthel.) Webb & Berthel. – NP – Macarones. (Neo – Cas)

Bulliarda vaillantii (Willd.) DC. – T scap – Medit.-Trop.

Crassula tillaea Lest.-Garl. – T scap – Medit.-Atl.

Kalanchoë daigremontiana Raym.-Hamet & H. Perrier – Ch succ – Africa (Neo – Cas)

Petrosedum sediforme (Jacq.) Grulich subsp. ***sediforme*** – Ch succ – Circum-Medit.

Phedimus stellatus (L.) Raf. – T scap – Circum-Medit.

Sedum album subsp. ***micranthum*** (Bast. ex DC.) Syme – Ch succ – Euro-Medit.

Sedum caeruleum L. – T scap – SW-Medit.

Sedum litoreum Guss. – T scap – Circum-Medit.

Sedum rubens L. – T scap – Euro-Medit.

Umbilicus horizontalis (Guss.) DC. – G bulb – Circum-Medit.

Umbilicus rupestris (Salisb.) Dandy – G rhiz – Medit.-Atl.

GERANIACEAE

- Erodium chium* (L.) Willd. – T scap – Circum-Medit.
Erodium ciconium (L.) L'Hér. – H bien – Euro-Medit.
Erodium cicutarium (L.) L'Hér. – H ros – Subcosmop.
Erodium malacoides (L.) L'Hér. subsp. *malacoides* – T scap – Medit.-Macarones.
Erodium moschatum (L.) L'Hér. – T scap – Circum-Medit.
Geranium molle L. – T scap – Subcosmop.
Geranium purpureum Vill. – T scap – Euro-Medit.
Geranium rotundifolium L. – T scap – Paleotemp.

LYTHRACEAE

- Punica granatum* L. – P caesp – E-Medit.-Turan. (Arch – Cas)

MYRTACEAE

- Eucalyptus camaldulensis* Dehnh. – P scap – Australia (Neo – Nat)
Eucalyptus globulus Labill. – P scap – Australia (Neo – Cas)
Myrtus communis L. – P caesp – Circum-Medit.

ZYGOPHYLLACEAE

- Tribulus terrestris* L. – T rept – Asiat. (Arch – Nat)

CELASTRACEAE

- Euonymus japonicus* Thunb. – P caesp – Asiat. (Neo – Cas)

CUCURBITACEAE

- Bryonia marmorata* E. Petit – G rhiz – Endem. SA-CO
Cucumis melo L. – T scap – Paleotrop. (Arch – Cas)
Cucumis sativus L. – T scap – Paleotrop. (Arch – Cas)
Cucurbita maxima Duchesne subsp. *maxima* – T scap – Neotrop. (Neo – Cas)
Cucurbita pepo L. subsp. *pepo* – T scap – Neotrop. (Neo – Cas)
Ecballium elaterium (L.) A.Rich. – G bulb – Euro-Medit.

FABACEAE

- Acacia saligna* (Labill.) H.L.Wendl. – P caesp – Australia (Neo – Inv)
Anagyris foetida L. – P caesp – S-Medit.
Astragalus hamosus L. – T scap – Medit.-Turan.
Astragalus pelecinus (L.) Barneby subsp. *pelecinus* – T scap – Circum-Medit.
Bituminaria bituminosa (L.) C.H.Stirt. – H scap – Euro-Medit.
Cytisus laniger DC. – P caesp – Circum-Medit.
Ceratonia siliqua L. – P scap – E-Medit. (Arch – Nat)
Ervilia hirsuta (L.) Opiz – T scap – Paleotemp.
Ervum gracile DC. – T scap – Euro-Medit.
Ervum pubescens DC. – T scap – Euro-Medit.
Genista corsica (Loisel.) DC. – NP – Endem. SA-CO
Lathyrus annuus L. – T scap – Euro-Medit.
Lathyrus cicera L. – T scap – Euro-Medit.

- Lathyrus clymenum* L. (sensu Gallego 1999) – T scap – Circum-Medit.
Lathyrus oleraceus subsp. *biflorus* (Raf.) H. Schaef., Coulot & Rabaute – T scap – Medit.-Turan.
Lathyrus oleraceus Lam. subsp. *oleraceus* – T scand – Circum-Medit. (Arch – Cas)
Lotus angustissimus L. – T scap – Euro-Medit.
Lotus cytisoides subsp. *conradiae* Gamisans – Ch suffr – Endem. SA-CO
Lotus edulis L. – T scap – Circum-Medit.
Lotus ornithopodioides L. – T scap – Circum-Medit.
Lotus parviflorus Desf. – T scap – Circum-Medit.
Lupinus angustifolius L. – T scap – Circum-Medit.
Lupinus gussoneanus J. Agardh. – T scap – Circum-Medit.
Medicago italica (Mill.) – T scap – W-Medit.
Medicago littoralis Loisel. – T scap – Euro-Medit.
Medicago marina L. – Ch rept – Medit.-Atl.
Medicago minima (L.) L. – T scap – Eurasiat.
Medicago orbicularis (L.) Bartal. – T scap – Euro-Medit.
Medicago polymorpha L. – T scap – Subcosmop.
Medicago praecox DC. – T scap – Circum-Medit.
Medicago rugosa Desr. – T scap – W-Medit.
Medicago truncatula Gaertn. – T scap – Circum-Medit.
Medicago turbinata (L.) All. – T scap – Circum-Medit.
Trigonella elegans (Salzm. ex Ser.) Coulot & Rabaute – T scap – S-Medit.
Trigonella italica (L.) Coulot & Rabaute – T scap – N-Medit.
Trigonella officinalis (L.) Coulot & Rabaute – H bien – Subcosmop.
Trigonella sicula (Turra) Coulot & Rabaute – T scap – S-Medit. (Arch – Inv)
Trigonella smalii Coulot & Rabaute – T scap – Subcosmop.
Trigonella sulcata (Desf.) Coulot & Rabaute – T scap – Circum-Medit.
Ononis diffusa Ten. – T scap – Circum-Medit.
Ononis natrix subsp. *ramosissima* (Desf.) Batt. – H suffr – Circum-Medit.
Ononis reclinata L. – T scap – Circum-Medit.
Ononis variegata L. – T scap – Circum-Medit.
Ornithopus compressus L. – T scap – Euro-Medit.
Ornithopus pinnatus (Mill.) Druce – T scap – Medit.-Atl.
Robinia pseudacacia L. – P caesp – Neotrop. (Neo – Nat)
Scorpiurus muricatus L. – T scap – Euro-Medit.
Spartium junceum L. – P caesp – Euro-Medit. (Arch – Cas)
Sulla coronaria (L.) Medik. – H scap – W-Medit. (Arch – Nat)
Trifolium angustifolium L. subsp. *angustifolium* – T scap – Euro-Medit.
Trifolium arvense L. subsp. *arvense* – T scap – Euro-Medit.-Irano-Turan.
Trifolium campestre Schreb. – T scap – W-Paleotemp.
Trifolium cherleri L. – T scap – Euro-Medit.
Trifolium glomeratum L. – T scap – Euro-Medit.
Trifolium scabrum L. – T scap – Euro-Medit.
Trifolium spumosum L. – T scap – Circum-Medit.
Trifolium stellatum L. – T scap – Euro-Medit.

- Trifolium subterraneum* L. subsp. ***subterraneum*** – T rept – Euro-Medit.
Trifolium suffocatum L. – T scap – Circum-Medit.
Trifolium tomentosum L. – T rept – W-Paleotemp.
Tripodion tetraphyllum (L.) Fourr. – T scap – Circum-Medit.
Vachellia karroo (Hayne) Banfi & Galasso – P caesp – S-Africa (Neo – Inv)
Vicia bithynica (L.) L. – T scap – Euro-Medit.
Vicia angustifolia L. – T scap – Subcosmop.
Vicia faba L. – T scap – S-Medit. (Arch – Cas)
Vicia pseudocracca Bertol. – T scap – Circum-Medit.
Vicia sativa L. – T scap – Medit.-Turan.
Vicia villosa Roth – T scap – Euro-medit.

CYTINACEAE

- Cytinus hypocistis* (L.) L. – G rad – Medit.-Macarones.

EUPHORBIACEAE

- Euphorbia characias* L. – NP – Circum-Medit.
Euphorbia dendroides L. – NP – C-Medit.
Euphorbia exigua L. subsp. ***exigua*** – T scap – Euro-Medit.
Euphorbia helioscopia L. subsp. ***helioscopia*** – T scap – Cosmop.
Euphorbia paralias L. – Ch frut – Medit.-Atl.
Euphorbia peplis L. – T rept – Euro-Medit.
Euphorbia peplus L. – T scap – Cosmop.
Euphorbia pithyusa L. subsp. ***pithyusa*** – Ch suffr – W-Medit.
Euphorbia pterococca Brot. – T scap – W-Medit.-Macarones.
Euphorbia segetalis L. – Ch suffr – W-Medit.
Euphorbia terracina L. – H scap – Circum-Medit.
Mercurialis annua L. – T scap – Paleotemp.
Ricinus communis L. – P scap – Paleotrop. (Arch – Inv)

LINACEAE

- Linum usitatissimum* subsp. ***angustifolium*** (Huds.) Thell. – H bienn – Euro-Medit.-Atl.
Linum corymbulosum Rchb. – T scap – Circum-Medit.
Linum strictum L. – T scap – Circum-Medit.
Linum tenuifolium L. – Ch suffr – S-Medit.
Linum trigynum L. – T scap – Euro-Medit.

SALICACEAE

- Populus alba* L. – P scap – Paleotemp.

OXALIDACEAE

- Oxalis corniculata* L. – H rept – Cosmop. (Arch D – Nat)
Oxalis pes-caprae L. – G bulb – S-Africa (Neo – Inv)
Oxalis articulata Savigny – G bulb – Neotrop. (Neo – Nat)

MORACEAE

Ficus carica L. – P scap – Medit.-Turan. (Arch D – Nat)

RHAMNACEAE

Rhamnus alaternus L. subsp. *alaternus* – P caesp – Circum-Medit.

ROSACEAE

Aphanes australis Rydb. – T scap – W-Medit. (Db)

Malus domestica (Borkh.) Borkh. – P scap – Euroasiat. (Arch – Cas)

Poterium sanguisorba subsp. *balearicum* (Bourg. ex Nyman) Stace – H scap – Medit.

Prunus dulcis (Mill.) D.A. Webb – P scap – Medit.-Turan. (Arch – Cas)

Pyrus spinosa Forssk. – P caesp – Circum-Medit.

Rubus ulmifolius Schott – NP – Euro-Medit.

URTICACEAE

Parietaria judaica L. – H scap – Euro-Medit.

Parietaria lusitanica L. subsp. *lusitanica* – T rept – Circum-Medit.

Urtica membranacea Poir. – T scap – S-Medit.

Urtica pilulifera L. – T scap – S-Medit.

Urtica urens L. – T scap – Boreo-Trop.

BRASSICACEAE

Arabidopsis thaliana (L.) Heynh. – T scap – Cosmop.

Biscutella morisiana Raffaelli – T scap – Endem. SA-CO

Brassica insularis Moris – Ch suffr – Endem. SA-CO-SI-TN

Brassica napus L. – H scap – Asiat. (Arch – Nat)

Brassica nigra (L.) W.D.J.Koch – T scap – Euro-Medit. (Arch D – Nat)

Brassica rapa subsp. *campestris* (L.) A.R.Clapham – H scap – Euro-Medit. (Arch – Cas)

Brassica tournefortii Gouan – T scap – S-Medit.

Bunias erucago L. – T scap – Euro-Medit.

Cakile maritima Scop. subsp. *maritima* – T scap – Circum-Medit.

Capsella bursa-pastoris (L.) Medik. subsp. *bursa-pastoris* – H bienn – Cosmop.

Cardamine hirsuta L. – T scap – Cosmop.

Diplotaxis viminea (L) DC. – T scap – W-Medit.-Atl.

Eruca vesicaria (L.) Cav. – T scap – Euro-Medit.-Irano-Turan. (Arch D – Nat)

Erysimum cheiri (L.) Crantz – Ch suffr – Euro-Medit. (Arch – Cas)

Hirschfeldia incana (L.) Lagr.-Foss. subsp. *incana* – H scap – W-Medit.

Lepidium graminifolium L. subsp. *graminifolium* – Ch suffr – W-Medit.

Lobularia maritima (L.) Desv. – H scap – Circum-Medit.

Marcus-kochia ramosissima (Desf.) Al-Shehbaz – T scap – W-Medit.

Matthiola tricuspidata (L.) R.Br. – T scap – Circum-Medit.

Raphanus raphanistrum L. subsp. *raphanistrum* – T scap – Euro- Medit.

Rapistrum rugosum (L.) All. – T scap – Euro-Medit.

Sinapis alba L. subsp. *alba* – T scap – E-Medit. (Arch – Nat)

Sinapis arvensis L. subsp. *arvensis* – T scap – Circum-Medit.

Sisymbrium erysimoides Desf. – T scap – S-Medit.

Sisymbrium officinale (L.) Scop. – T scap – Subcosmop.

Succowia balearica (L.) Medik. – T scap – W-Medit.

Teesdalia coronopifolia (J.P. Bergeret) Thell. – T scap – Euro-Medit.

RESEDACEAE

Reseda alba L. – H scap – Circum-Medit.

Reseda luteola L. – H scap – Circumbor.

CISTACEAE

Cistus creticus subsp. *eriocephalus* (Viv.) Greuter & Burdet – NP – Circum-Medit.

Cistus monspeliensis L. – NP – Medit.-Macarones.

Cistus salviifolius L. – NP – Circum-Medit.

Fumana thymifolia (L.) Spach. ex Webb – Ch suffr – Circum-Medit.

Helianthemum salicifolium (L.) Mill. – T scap – Euro-Medit.

Tuberaria guttata (L.) Fourr. – T scap – Euro-Medit.

MALVACEAE

Alcea rosea L. – H scap – E-Medit. (Arch – Cas).

Malva arborea (L.) Webb & Berthel. – H bienn – Euro-Medit.

Malva multiflora (Cav.) Soldano, Banfi & Galasso – T scap – Circum-Medit.

Malva nicaeensis All. – T scap – Circum-Medit.

Malva olbia (L.) Alef. – P caesp – Circum-Medit.

Malva parviflora L. – T scap – Euro-Medit.

Malva sylvestris L. – H scap – Subcosmop.

THYMELAEACEAE

Daphne gnidium L. – P caesp – Medit.-Macarones.

Thymelaea hirsuta (L.) Endl. – NP – Circum-Medit.

Thymelaea tartonraira (L.) All. subsp. *tartonraira* – NP – Circum-Medit.

ANACARDIACEAE

Pistacia lentiscus L. – P caesp – Circum-Medit.

Schinus molle L. – P scap – Neotrop. (Neo – Cas).

RUTACEAE

Ruta chalepensis L. – Ch suffr – S-Medit.

SIMAROUBACEAE

Ailanthus altissima Mill. – P scap – Asiat. (Neo – Cas).

ERICACEAE

Erica arborea L. – P caesp – Circum-Medit.

Erica scoparia L. subsp. *scoparia* – P caesp – Circum-Medit.

PRIMULACEAE

- Lysimachia arvensis* subsp. *latifolia* (L.) Peruzzi – T rept – Euro-Medit.
Lysimachia foemina (Mill.) U. Manns & Anderb. – T rept – Subcosmop.
Lysimachia linum-stellatum L. – T scap – Circum-Medit.
Lysimachia nardii Arrigoni – T rept – Circum-Medit. (Arch – Nat).
Samolus valerandi L. – H scap – Subcosmop.

BORAGINACEAE

- Borago officinalis* L. – T scap – Euro-Medit.
Cynoglossum creticum Mill. – H bienn – Euro-Medit.
Echium creticum L. subsp. *creticum* – H bien – W-Medit.
Echium italicum L. – H bienn – Euro-Medit.
Echium plantagineum L. – H bienn – Euro-Medit.
Echium vulgare L. – H bienn – Europ.
Heliotropium europaeum L. – T scap – Euro-Medit.-Turan.
Myosotis arvensis (L.) Hill subsp. *arvensis* – T scap – Euroasiat.
Myosotis ramosissima Rochel ex Schult. subsp. *ramosissima* – T scap – Euroasiat.

APOCYNACEAE

- Nerium oleander* L. subsp. *oleander* – P caesp – S-Medit.
Vincetoxicum hirundinaria Medik. subsp. *hirundinaria* – H scap – Euroasiat.

GENTIANACEAE

- Centaurium erythraea* Raf. subsp. *erythraea* – T scap – Paleotemp.
Centaurium maritimum (L.) Fritsch – T scap – W-Medit.
Centaurium tenuiflorum (Hoffmanns. & Link) Fritsch subsp. *tenuiflorum* – T scap –
 Medit.-Atl.
Schenkia spicata (L.) G.Mans. – T scap – Euro-Medit.

RUBIACEAE

- Crucianella maritima* L. – Ch suffr – Circum-Medit.
Galium aparine L. – T scap – Eurasiat.
Galium murale (L.) All. – T scap – Circum-Medit.
Galium spurium L. – T scap – Euro-Medit.-Irano-Turan.
Galium verrucosum Huds. subsp. *verrucosum* – T scap – Circum-Medit.
Rubia peregrina L. – P lian – Circum-Medit.
Sherardia arvensis L. – T scap – Subcosmop.
Thelygonum cynocrambe L. – T scap – Circum-Medit.
Valantia muralis L. – T scap – Circum-Medit.

ACANTHACEAE

- Acanthus mollis* L. subsp. *mollis* – H scap – W-Medit. (Arch – Cas).

BIGNONIACEAE

- Tecomaria capensis* (Thunb.) Spach – T scap – S-Africa (Neo – Cas).

LAMIACEAE

- Ajuga iva* (L.) Schreber subsp. *iva* – Ch suffr – Circum-Medit.
Clinopodium vulgare subsp. *arundanum* (Boiss.) Nyman – H scap – Circumbor.
Lamium amplexicaule L. – T scap – Paleotemp.
Lavandula stoechas L. subsp. *stoechas* – NP – Circum-Medit.
Marrubium vulgare L. – H scap – Subcosmop.
Melissa officinalis subsp. *altissima* (Sm.) Arcang. – H scap – Euro-Medit.
Mentha pulegium L. subsp. *pulegium* – H scap – Subcosmop.
Micromeria graeca (L.) Benth. ex Rchb. subsp. *graeca* – Ch suffr – Circum-Medit.
Salvia rosmarinus Spenn. – NP – Circum-Medit.
Salvia verbenaca L. – H scap – Medit.-Atl.
Stachys major (L.) Bartolucci & Peruzzi – Ch frut – Circum-Medit.
Stachys romana (L.) E.H.L. Krause – T scap – Circum-Medit.
Teucrium marum L. – Ch suffr – Endem. SA-CO-BL-AT-H
Vitex agnus-castus L. – P caesp – Circum-Medit.

OLEACEAE

- Olea europaea* var. *sylvestris* Brot. – P scap – Circum-Medit.
Phillyrea angustifolia L. – P caesp – W-Medit.
Phillyrea latifolia L. – P scap – Circum-Medit.
Phillyrea latifolia var. *media* L. C.K. Schneid – P scap – Circum-Medit.

OROBANCHACEAE

- Bellardia trixago* (L.) All. – T scap – Euro-Medit.
Bellardia viscosa (L.) Fisch. & C.A. Mey. – T scap – Medit.-Atl.
Orobanche amethystea Thuill. – T par – Euro-Medit.
Orobanche minor Sm. – T par – Subcosmop.
Phelipanche ramosa (L.) Pomel – T par – Circum-Medit.

PLANTAGINACEAE

- Callitriche palustris* L. – I rad – Circumbor.
Cymbalaria aequitriloba (Viv.) A.Chev. subsp. *aequitriloba* – Ch rept – Endem. SA-CO-BL-AT
Linaria arvensis (L.) Desf. – T scap – Medit.-Atl.
Linaria pelisseriana (L.) Mill. – T scap – Medit.-Atl.
Linaria simplex (Willd.) Desf. – T scap – Euro-Medit.
Misopates orontium (L.) Raf. – T scap – Paleotemp.
Plantago afra L. – T scap – Euro-Medit.
Plantago bellardii All. subsp. *bellardii* – T scap – S-Medit.
Plantago coronopus L. – T scap – Euro-Medit.
Plantago crassifolia Forssk. – H ros – Circum-Medit.
Plantago lagopus L. – T scap – Circum-Medit.
Plantago lanceolata L. – H ros – Cosmop.
Plantago macrorrhiza Poir. – H ros – W-Medit.
Plantago major L. – H ros – Euroasiat.

Plantago weldenii Rchb. – T scap – Euro-Medit.

Veronica cymbalaria Bodard subsp. *cymbalaria* – T scap – Euro-Medit.

Veronica persica Poir. – T scap – Irano-Turan. (Neo – Nat).

SCROPHULARIACEAE

Myoporum insulare R. Br. – P caesp – Australia (Neo – Nat).

Myoporum tetrandrum (Labill.) Domin – P caesp – Australia (Neo – Nat).

Scrophularia trifoliata L. – H scap – Endem. SA-CO-AT

Verbascum conocarpum Moris subsp. *conocarpum* – H bienn – Endem. SA-CO-AT

Verbascum sinuatum L. – H bienn – Euro-Medit.

VERBENACEAE

Lantana camara L. – NP – Neotrop. (Neo – Cas).

Verbena officinalis L. – H scap – Neotrop.

CONVOLVULACEAE

Convolvulus althaeoides L. – H scand – W-Medit.

Convolvulus arvensis L. – G rhiz – Cosmop.

Convolvulus sepium L. – H scand – Paleotemp.

Convolvulus sicularis subsp. *elongatus* Batt. – T scap – S-Medit.

Convolvulus soldanella L. – G rhiz – Cosmop.

Cuscuta epithymum (L.) L. subsp. *kotschy* (Des Moul.) Arcang. – T par – Circum-Medit.

SOLANACEAE

Datura innoxia Mill. – T scap – Neotrop. (Neo – Nat)

Datura stramonium L. – T scap – Neotrop. (Neo – Nat)

Datura wrightii Regel – T scap – Neotrop. (Neo – Nat)

Hyoscyamus albus L. – H bienn – Euro-Medit.

Nicotiana glauca Graham – NP – Neotrop. (Neo – Inv)

Solanum dulcamara L. – NP – Paleotemp.

Solanum linnaeanum Hepper & P.-M. L. Jaeger – NP – S-Africa (Neo – Nat)

Solanum lycopersicum L. – T scap – Neotrop. (Neo – Cas)

Solanum nigrum L. – T scap – Cosmop.

APIACEAE

Ammi majus L. – T scap – Euro-Medit.-Irano-Turan.

Ammoides pusilla (Brot.) Breistr. – T scap – Circum-Medit.

Bupleurum odontites L. – T scap – Medit.-Irano-Turan.

Bupleurum semicompositum L. – T scap – Circum-Medit.

Bupleurum tenuissimum L. – T scap – Euro-Medit.-Irano-Turan.

Crithmum maritimum L. – Ch suffr – Euro-Medit.

Daucus carota L. subsp. *carota* – H scap – Cosmop.

Daucus carota subsp. *hispanicus* (Gouan) Thell. – H bien – W-Medit.

Daucus carota subsp. *maritimus* (Lam.) Batt. – H bien – W-Medit.

Daucus carota subsp. *maximus* (Desf.) Ball – H bien – Euro-Medit.

Daucus pumilus (L.) Hoffmanns. & Link – T scap – Circum-Medit.
Echinophora spinosa L. – H scap – Euro-Medit.
Eryngium campestre L. – H scap – Euro-Medit.
Eryngium maritimum L. – G rhiz – Medit.-Atl.
Ferula arrigonii Bocchieri – H scap – Endem. SA-CO
Ferula communis L. subsp. *communis* – H scap – S-Medit.
Foeniculum vulgare Mill. – H scap – S-Medit.
Scandix pecten-veneris L. subsp. *pecten-veneris* – T scap – Subcosmop.
Smyrniolum olusatrum L. – H bienn – Medit.-Atl.
Thapsia garganica L. subsp. *garganica* – H scap – S-Medit.
Tordylium apulum L. – T scap – Circum-Medit.
Torilis nodosa (L.) Gaertn. subsp. *nodosa* – T scap – Euro-Medit.-Turan.

ASTERACEAE

Achillea maritima (L.) Ehrend. & Y.P.Guo subsp. *maritima* – Ch suffr – Medit.-Atl.
Andryala integrifolia L. – T scap – Euro-Medit.
Anthemis arvensis L. – T scap – Subcosmop.
Artemisia arborescens (Vaill.) L. – NP – SW-Medit.
Bellis annua L. subsp. *annua* – T scap – Medit.-Macarones.
Bellis perennis L. – H ros – Circumbor.
Calendula arvensis (Vaill.) L. – T scap – Euro-Medit.
Calendula officinalis L. – T scap – Medit.-Turan. (Arch – Cas)
Carduus fasciculiflorus Viv. – H bien – Endem. SA-CO-AT
Carduus pycnocephalus L. subsp. *pycnocephalus* – H bienn – Euro-Medit.-Turan.
Carlina corymbosa L. – H scap – Circum-Medit.
Carlina gummifera (L.) Less. – H ros – S-Medit.
Carlina lanata L. – T scap – Circum-Medit.
Carthamus lanatus L. – T scap – Euro-Medit.
Centaurea calcitrapa L. – H bien – Medit.-Atl.
Centaurea napifolia L. – T scap – SW-Medit.
Centaurea sphaerocephala L. subsp. *sphaerocephala* – H scap – Circum-Medit.
Chamaemelum fuscatum (Brot.) Vasc. – T scap – W-Medit.
Chondrilla juncea L. – H scap – Euro-Medit.
Cichorium intybus L. – H scap – Cosmop.
Cirsium vulgare (Savi) Ten. subsp. *sylvaticum* (Tausch) Arènes – H bien – Subcosmop.
Cladanthus mixtus (L.) Chevall. – T scap – Circum-Medit.
Crepis bellidifolia Loisel. – T scap – W-Medit.
Crepis hyemalis (Biv.) Ces., Pass. & Gibelli – T scap – Endem. SA-SI
Crepis vesicaria L. subsp. *vesicaria* – T scap – Medit.-Atl.
Crupina crupinastrum (Moris) Vis. – T scap – Circum-Medit.
Cynara cardunculus L. subsp. *cardunculus* – H scap – Circum-Medit.

- Dimorphotheca ecklonis* DC. – Ch scap – S-Africa (Neo – Cas)
Dittrichia graveolens (L.) Greuter – T scap – Medit.-Turan.
Dittrichia viscosa (L.) Greuter subsp. *viscosa* – H scap – Euro-Medit.
Erigeron bonariensis L. – T scap – Neotrop. (Neo – Inv)
Erigeron canadensis L. – T scap – Neotrop. (Neo – Inv)
Erigeron sumatrensis Retz. – T scap – Neotrop. (Neo – Inv)
Eriocephalus africanus L. – P – S Africa (Neo – Cas).
Filago asterisciflora (Lam.) Sweet – T rept – Circum-Medit.
Filago germanica (L.) Huds. – T scap – Paleotemp.
Filago pygmaea L. – T rept – Circum-Medit.
Filago pyramidata L. – T scap – Euro-Medit.
Galactites tomentosus Moench – H bienn – Circum-Medit.
Galatella pannonica (Jacq.) Galasso, Bartolucci & Ardenghi – H bien – Euro-Medit.-
 Irano-Turan.
Glebionis coronaria (L.) Spach. – T scap – Circum-Medit. (Arch – Inv)
Hedypnois rhagadioloides (L.) F.W.Schmidt – T scap – Circum-Medit.
Helianthus annuus L. subsp. *annuus* – T scap – Neotrop. (Neo – Cas)
Helichrysum italicum subsp. *tyrrhenicum* (Bacch., Brullo & Giusso) Herrando,
 J.M.Blanco, L.Sáez & Galbany – Ch suffr – Endem. SA-CO
Helminthotheca echioides (L.) Holub – T scap – Euro-Medit.
Hyoseris scabra L. – T ros – Circum-Medit.
Hypochaeris achyrophorus L. – T scap – Circum-Medit.
Hypochaeris glabra L. – T scap – Euro-Medit.
Hypochaeris radicata L. – T scap – Circum-Medit.
Lactuca sativa L. subsp. *sativa* – H bienn – Medit. (Arch – Cas)
Lactuca sativa subsp. *serriola* (L.) Galasso, Banfi, Bartolucci & Ardenghi – T scap –
 Euro-Medit.
Leontodon tuberosus L. – H ros – Circum-Medit.
Limbarda crithmoides subsp. *longifolia* (Arcang.) Greuter – Ch suffr – Circum-Medit.
Logfia gallica (L.) Cosson & Germ. – T scap – Euro-Medit.
Onopordum illyricum L. – H bien – NE-Medit.
Pallenis spinosa (L.) Cass. subsp. *spinosa* – H bienn – Euro-Medit.
Phagnalon rupestre subsp. *illyricum* (H.Lindb.) Ginzb. – Ch suffr – W-Medit.
Phagnalon saxatile (L.) Cass. – Ch suffr – W-Medit.
Pulicaria odora (L.) Rchb. – H scap – Euro-Medit.
Pulicaria sicula (L.) Moris – T scap – Circum-Medit.
Reichardia picroides (L.) Roth – H scap – Circum-Medit.
Rhagadiolus stellatus (L.) Gaertn. – T scap – Euro-Medit.
Scolymus hispanicus L. – H bienn – Euro-Medit.
Scolymus maculatus L. – T scap – S-Medit.
Senecio leucanthemifolius Poir. subsp. *leucanthemifolius* – T scap – Circum-Medit.
Senecio lividus L. – T scap – Circum-Medit.
Senecio transiens (Rouy) Jeanm. – T scap – Endem. SA-CO
Senecio vulgaris L. subsp. *vulgaris* – T scap – Cosmop.
Silybum marianum (L.) Gaertn. – H bienn – Medit.-Turan.

Sonchus asper (L.) Hill subsp. *asper* – T scap – Subcosmop.

Sonchus bulbosus (L.) N. Kilian & Greuter subsp. *bulbosus* – G bulb – Circum-Medit.

Sonchus oleraceus L. – T scap – Subcosmop.

Sonchus tenerrimus L. – T scap – Circum-Medit.

Symphytotrichum squamatum (Spreng.) G.L.Nesom – T scap – Neotrop. (Neo – Inv).

Tolpis umbellata Bertol. – T scap – Circum-Medit.

Tolpis virgata (Desf.) Bertol. subsp. *virgata* – H scap – Circum-Medit.

Urospermum dalechampii (L.) F.W. Schmidt – H scap – Euro-Medit.

Urospermum picroides (L.) Scop. ex F.W. Schmidt – T scap – Euro-Medit.

Xanthium italicum Moretti – T scap – Neotrop. (Neo – Nat).

Xanthium spinosum L. – T scap – Neotrop. (Neo – Inv).

CAMPANULACEAE

Campanula erinus L. – T scap – Circum-Medit.

Jasione montana L. – H bienn – Euro-Medit.

DIPSACACEAE

Dipsacus ferox Loisel. – H bienn – Endem. SA-CO

Lomelosia rutifolia (Vahl) Avino & P.Caputo – H scap – W-Medit.

Sixalis atropurpurea (L.) Greuter & Burdet – H bien – Circum-Medit.

VALERIANACEAE

Centranthus calcitrapae (L.) Dufr. subsp. *calcitrapae* – T scap – Circum-Medit.

Valerianella dentata (L.) Pollich – T scap – Medit.-Atl.

Valerianella microcarpa Loisel. – T scap – Circum-Medit.

FAGACEAE

Quercus calliprinos Webb. – P scap – E-Medit.

Quercus ilex L. – P scap – Circum-Medit.

Quercus suber L. – P scap – Circum-Medit.

Results

The inventory of the vascular flora consists of 653 taxa, 84% of which are native (5% of them being endemic) and 16% non-native (10% neophytes, 6% archaeophytes) (Fig. 2), including 500 species, 148 subspecies, three varieties and two hybrids, from 89 families and 374 genera.

The native component includes 406 species, 135 subspecies, two variety and one hybrid, belonging to 75 families and 313 genera. The best represented phylum (Table 2) is *Magnoliophyta* (532 taxa), in which *Eudicotyledones* represent the largest systematic group, with 402 taxa, followed by 128 *Monocotyledones* taxa. The phylum *Pteridophyta* is represented by nine taxa, *Pinophyta* only by three taxa.

Among alien taxa, the best represented phylum (Table 2) is *Magnoliophyta* (107 taxa), in which *Eudicotyledones* represent the largest systematic group with 79 taxa, followed by *Monocotyledones* (28 taxa) and *Pinophyta* (two taxa).




Fig. 2. The total vascular flora of the MPACC.

Among the native flora, the best represented family is *Asteraceae* (68 taxa) followed by *Poaceae* (60 taxa) and *Fabaceae* (59 taxa; Fig. 3).

Life-form analysis (Fig. 4) of native species revealed dominance of therophytes (49%) followed by hemicryptophytes (21%), geophytes (14%) and chamaephytes (7%). Phanerophytes are 5%, nano-phanerophytes 3%, hydrophytes only 1%.

As concerns chorology of native taxa (Fig. 5), the Mediterranean element is largely prevailing (81%, 440 taxa), followed by the Subcosmopolitan (6%, 33 taxa), Paleotemperate (4%, 22 taxa) and Cosmopolitan (3%, 17 taxa) elements.

In the chorological spectrum of the native Mediterranean species (Fig. 6a) most represented are the Circum-Mediterranean (37%, 162 taxa) and Euro-Mediterranean (27%, 119 taxa), followed by W-Mediterranean (9%, 40 taxa) and Endemic (6%, 29 taxa).

Table 2. Taxa numbers and percentages per phylum in the native and alien vascular flora of the MPACC.

Phylum	Native	Alien	Native	Alien
	N°	N°	%	%
Magnoliophyta - Eudicotyledones	402	79	73,90	72,48
Magnoliophyta - Monocotyledones	128	28	23,53	25,69
Magnoliophyta - Magnolides	2	0	0,37	0
Pinophyta	3	2	0,55	1,83
Pteridophyta	9	0	1,65	0
Total	544	109	100	100




Fig. 3. Taxa numbers in the main families of the native vascular flora of the MPACC.




Fig. 4. Comparison of taxa numbers per life-form in the native and alien vascular flora of the MPACC.




Fig. 5. Taxa numbers in the chorological types of the native vascular flora of the MPACC.

Among endemic plants (Fig. 6b), Sardo-Corsican taxa are predominant (55%, 16 taxa), followed by Sardo-Corsican-Tuscan Archipelago (14%, four taxa) and Sardinian taxa (10%, three taxa).

The non-native component includes 94 species, 13 subspecies, one variety and one hybrid, belonging to 40 families and 81 genera. Of the 109 alien taxa (Fig. 7), 25 (23%) are invasive, 40 (37%) naturalized and 44 (40%) casual. Among neophytes (67 taxa) the number of invasive taxa is 21 (19%), naturalized taxa are 20 (18%) and casual 26 (24%). Archaeophytes (42 taxa) have four invasive taxa (4%), 20 naturalized (19%) and 18 casual taxa (16%).

In the alien flora, the most represented family is *Asteraceae* (12 taxa) followed by *Fabaceae* (nine taxa) and *Poaceae* (nine taxa) (Fig. 8).

The best represented life-forms (Fig. 4) among alien taxa are therophytes (33%) and phanerophytes (32%), followed by geophytes (13%), hemicryptophytes (11%), chamaephytes (6%) and nano-phanerophytes (5%). There are no alien hydrophytes.

As concerns the geographic origin of alien species (Fig. 9), the American element (35%, 38 taxa) prevails, followed by the Mediterranean (27%, 30 taxa), the South African (13%, 14 taxa) and Asian (9%, 10 taxa) elements.





Fig. 6. Chorological spectrum of the Mediterranean (a) and Endemic (b) component of the native vascular flora of the MPACC.




Fig. 7. Number of taxa per diffusion status among archaeophytes (Arch) and neophytes (Neo) of the MPACC.




Fig. 8. Taxa numbers in the main families of the alien vascular flora of the MPACC.




Fig. 9. Number of alien vascular taxa of the MPACC grouped by geographic origin.

Discussion

The floristic composition of the territory is similar to that of other Sardinian territories, especially as regards native families (e.g. Bacchetta 2006). Predominant are *Asteraceae*, *Poaceae* and *Fabaceae* which make up more than a third (34%) of the species in the MPACC's flora (together with *Caryophyllaceae*, *Apiaceae* and *Brassicaceae* almost 50%). As regards the alien component, after the three families that have the greatest number of taxa (*Asteraceae*, *Poaceae* and *Fabaceae*), there are also families such as *Aizoaceae* and *Cactaceae* that are exclusively represented by alien species.

The life-form spectrum of the MPACC's flora is consistent with those of other floras of neighbouring territories and highlights the Mediterranean nature of the climate in which therophytes (49%) and hemicryptophytes (21%) have a clear dominance over perennial biological forms (Bocchieri & al. 2008). The high percentage of geophytes (14%) appears to be linked to the anthropogenic use of the territory, in particular to the practice of fires and to agro-pastoral activities, but also proves the abundance of psammophilous habitats in which geophytes are important for the creation of dunes. The low percentage of phanerophytes (5%) and nano-phanerophytes (3%) reflects the low degree of woody cover.

The low percentage of hydrophytes is related to the almost absence of freshwater habitats, the five species present being related to sea, or pond water. As regards the alien component, the greater percentage of phanerophytes (32%) is emphasized, which, together with nano-phanerophytes (5%) become the prevalent biological form, testifying the use of species mainly introduced for reforestation, forestation and ornamental use, as already found for the alien vascular flora of Sardinia (Puddu & al. 2016).

Our analyses highlight the high floristic diversity of the investigated area, as testified by the presence of plants of high phytogeographical value, including 29 endemic species mainly confined in the two islands. Among the endemics, three species are exclusive to Sardinia: *Aristolochia tyrrhena* E. Nardi & Arrigoni, *Limonium retirameum* Greuter & Burdet subsp. *retirameum* and *Silene valsecchiai* Bocchieri. The latter two are also exclusive to eastern Sardinia. Most of the endemics (16 taxa) consist of Sardo-Corsican elements: *Aristolochia rotunda* subsp. *insularis* (E.Nardi & Arrigoni) Gamisans; *Biscutella morisiana* Raffaelli; *Bryonia marmorata* E. Petit; *Dipsacus ferox* Loisel.; *Ferula arrigonii* Bocchieri; *Genista corsica* (Loisel) DC.; *Helichrysum italicum* subsp. *tyrrhenicum* (Bacch., Brullo & Giusso) Herrando, J. M. Blanco, L. Sáez & Galbany; *Lotus cytisoides* subsp. *conradiae* Gamisans; *Narcissus supramontanus* Arrigoni subsp. *cunicularium* Arrigoni; *Ophrys conradiae* Melki & Deschâtres; *O. exaltata* Ten. subsp. *morisii* (Martelli) Del Prete; *Portulaca sardoa* Danin, Bagella & Marrosu; *Prospero corsicum* (Boullu) J.-M. Tison; *Romulea requienii* Parl.; *Senecio transiens* (Rouy) Jeanm.; *Silene succulenta* subsp. *corsica* (DC.) Nyman. *S. valsecchiai* and *F. arrigonii* have the *locus classicus* on the island of Serpentara. The endemics shared with Corsica and the Tuscan Archipelago are *Arum pictum* L. f. subsp. *pictum*, *Carduus fasciculiflorus* Viv., *Scrophularia trifoliata* L. and *Verbascum conocarpum* Moris subsp. *conocarpum*. Noteworthy is the abundance of the endemic *Brassica insularis* Moris, listed in Annexes II and IV of the Habitats Directive 92/43/EEC. This species, considered a Tyrrhenian endemic, is found only on the island of Cavoli, where one of the most important populations of Sardinia is present, the island taking its name from this species. The island of Serpentara is also probably a phytotoponym, which seems to take its name from *Helicodiceros muscivorus* (L. f.) Engl. (Biondi & al. 1993), endemic to Sardinia, Corsica, and the Balearic Islands.

Like most of the coastal territories of the Mediterranean area, also the study area is not immune from the spread of alien plant species, which make up 17% of the total flora. In particular, the high presence of invasive species is alarming, above all because they are a threat to the biodiversity of the most sensitive habitats, such as those of coastal dunes (Podda & al. 2018; Mayoral & al. 2020). In fact, among the 25 invasive species, the most common are neophytes introduced in gardens and public greenery as ornamental species, which escaped cultivation and become invasive in coastal habitats, such as *Carpobrotus acinaciformis* (L.) L. Bolus, *Carpobrotus edulis* (L.) N. E. Br., *Malephora crocea* (Jacq.) Schwantes and *Mesembryanthemum cordifolium* L.f., all belonging to the South African family *Aizoaceae*. In addition, several American plant species belonging to the *Agave* genus [*Agave americana* L., *Agave fourcroydes* Lem., *Agave ingens* Brg. var. *picta* (Salm.) Bgr., *Agave salmiana* subsp. *ferox* (K.Koch) Hochstätter], two *Fabaceae* [*Acacia saligna* (Labill.) H. L. Wendl. and *Vachellia karroo* (Hayne) Banfi & Galasso] and two *Cactaceae* [*Opuntia ficus-indica* (L.) Mill. e *Austrocylindropuntia subulata* (Mühlenpf.) Backeb] occur. There are also other species which are considered invasive in other territories, that have not yet shown their potential, but which could become so in the future such as *Ailanthus altissima* (Mill.) Swingle, *Cortaderia selloana* (Schult.) Asch. & Graebn. and *Cenchrus longisetus* M.C. Johnst. Differently, the archaeophytes of the study area are naturalized species, integrated with the flora, that show very little invasiveness since only four taxa [*Arundo donax* L., *Glebionis coronaria* (L.) Spach, *Ricinus communis* L., *Trigonella sicula* (Turra) Coulot & Rabaute] can be considered as invasive, especially in wetlands and synanthropic habitats.

Acknowledgements

This research was partially supported by LIFE13 NAT/IT/000433 RES MARIS project (2014-2018) and finalized under the RES MARIS post-LIFE.

References

- AA.VV. 2014: Layman's Report PROVIDUNE. Conservazione e ripristino di habitat dunali nei siti delle Province di Cagliari, Caserta e Matera. – LIFE07NAT/IT/000519 PROVIDUNE.
- Acunto, S., Bacchetta, G., Bordigoni, A., Cadoni, N., Cinti, M. F., Duràn Navarro, M., Frau, F., Lentini, L., Liggi, M. G., Masala, V., Meloni, F., Pinna, R., Podda, L. & Sanna, A. 2017: The LIFE+ project “RES MARIS - Recovering Endangered habitatS in the Capo Carbonara MARine area, Sardinia”: first results. – *Pl. Sociol.* **54**: 85-95.
- Angiosperm Phylogeny Group 2016: An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. – *Bot. J. Linn. Soc.* **181**: 1-20. <https://doi.org/10.1111/boj.12385>
- Arrigoni, P. V. 2006 - 2015: Flora dell'Isola di Sardegna, **1-6**. – Sassari.
- & Bocchieri, E. 1995: Caratteri fitogeografici della flora delle piccole isole circumsarde. – *Biogeographia* **18**: 63-90.
- & Di Tommaso, P. L. 1991: La vegetazione delle montagne calcaree della Sardegna centro-orientale. – *Boll. Soc. Sarda Sci. Nat.* **28**: 201-310.
- Bacchetta, G. 2006: Flora vascolare del Sulcis (Sardegna Sud-Occidentale, Italia). – *Guineana* **12**: 1-369.
- & Pontecorvo, C. 2005: Contribution to the knowledge of the endemic vascular flora of Iglesiente (SW Sardinia-Italy). – *Candollea* **60**: 481-501.
- , Farris, E. & Pontecorvo, C. 2012: A new method to set conservation priorities in biodiversity hotspots. – *Pl. Biosyst.* **146**: 638-648. <https://doi.org/10.1080/11263504.2011.642417>
- , Iiriti, G. & Mossa, L. 2005: La flora endemica del Sarrabus-Gerrei: un patrimonio da tutelare e gestire. Atti del Convegno “Analisi e sistemi di gestione del territorio (Sarrabus-Gerrei)”. – *Sinnai* 30/31.1.2004: 105-112.
- , Fenu, G., Mattana, E., Meloni, F. & Podda, L. 2007: Conservazione *ex situ* e *in situ* della biodiversità vegetale dell'Area Marina Protetta di Capo Carbonara (Sardegna sud-orientale). – *Fitosociologia* **44**: 157-164.
- , Bagella, S., Biondi, E., Farris, E., Filigheddu, R. & Mossa, L. 2009: Vegetazione forestale e serie di vegetazione della Sardegna (con rappresentazione cartografica alla scala 1:350.000). – *Fitosociologia* **46**: 3-82.
- , Duràn Navarro, M., Fenu, G., Iiriti, G., Meloni, F. & Podda, L. 2015: Contribution to the knowledge of the vascular flora of the Marine Protected Area Capo Carbonara (South East Sardinia). — In: Peruzzi, L. & Domina, G. (eds) Approfondimenti floristici e sistematici sulla flora d'Italia. Comunicazioni. Società Botanica Italiana, Gruppo per la Floristica, Sistematica ed Evoluzione. – Orto botanico di Roma, La Sapienza Università di Roma, 20-21 novembre 2015.
- , Fenu, G., Iiriti, G., Mattana, E., Meloni, F., Mulè, P. & Podda, L. 2006: Territory defence throughout conservation of the plant diversity: the project of the Protected Sea Area of Capo Carbonara (South Eastern Sardinia). – Pp. 302-307 in Acts of the 1st International Symposium on Environment, Identities and Mediterranean area (ISEIM 2006). – Corte, Ajaccio (France).
- , —, Guarino, R., Mandis, G., Mattana, E., Nieddu, G. & Scudu, C. 2013: Floristic traits and biogeographic characterization of the Gennargentu massif (Sardinia). – *Candollea* **68**: 209-220.

- , Bordigoni, A., Cinti, M. F., Frau, F., Lentini, L., Liggi, M. G., Meloni, F., Orrù, M., Podda, L. & Sanna, A. 2018: Manuale di buone pratiche e linee guida per la corretta fruizione e gestione degli habitat naturali del sistema spiaggia. Progetto LIFE+ RES MARIS. — Cagliari.
- Ballero, M. 1982: Etnoiatria nella zona di Villasimius (Sardegna sud-orientale). — Rend. Sem. Fac. Sci. Univ. Cagliari **52**: 115-125.
- Bartolucci, F., Peruzzi, L., Galasso, G., Albano, A., Alessandrini, A., Ardenghi, N. M. G., Astuti, G., Bacchetta, G., Ballelli, S., Banfi, E., Barberis, G., Bernardo, L., Bouvet, D., Bovio, M., Cecchi, L., Di Pietro, R., Domina, G., Fascetti, S., Fenu, G., Festi, F., Foggi, B., Gallo, L., Gottschlich, G., Gubellini, L., Iamónico, D., Iberite, M., Jiménez-Mejías, P., Lattanzi, E., Marchetti, D., Martinetto, E., Masin, R. R., Medagli, P., Passalacqua, N. G., Peccenini, S., Pennesi, R., Pierini, B., Poldini, L., Prosser, F., Raimondo, F. M., Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A., Scortegagna, S., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagensommer, R. P., Wilhalm, T. & Conti, F. 2018: An updated checklist of the vascular flora native to Italy. — Pl. Biosyst. **152**: 179-303. <https://doi.org/10.1080/11263504.2017.1419996>
- Biondi, E., Bocchieri, E., Brugiapaglia, E. & Mulas, B. 1993: La vegetazione dell'isola di Serpentara (Sardegna sud-orientale). — Boll. Soc. Sarda Sci. Nat. **29**: 115-130.
- Bocchieri, E. 1988: *Silene valsecchiaae* e *Ferula arrigonii*, due specie nuove della Sardegna. — Boll. Soc. Sarda Sci. Nat. **26**: 305-310.
- 1989: The flora of Serpentara island (Southern Sardinia): phytogeographic relevance and conservation requirements. — Coll. Phytosoc. **29**: 233-250.
- 2001: Endemismi e rarità tra la flora delle piccole isole della provincia di Cagliari (Sardegna). — Biogeographia **22**: 139-168.
- & Iriti, G. 2007a: L'esplorazione botanica nei territori del Sarrabus-Gerrei (Sardegna sud orientale). — Rend. Sem. Fac. Sci. Nat. Univ. Cagliari **77**: 85-99.
- & — 2007b: Nuovi dati sulla presenza di habitat e specie vegetali di interesse comunitario in alcuni Siti d'Importanza Comunitaria del Sarrabus-Gerrei (Sardegna sud orientale). — Fitosociologia **44**: 207-211.
- & — 2008: Valutazione dello stato di conservazione e monitoraggio dei principali sistemi psamofili del settore sud orientale della Sardegna. — Pp. 177-184 in: Atti Secondo Simposio Internazionale "Il monitoraggio costiero mediterraneo: problematiche e tecniche di misura". — Napoli.
- , —, & Pontecorvo, C. 2008: Flora vascolare del Capo di Monte Santu (Sardegna centro orientale). — Webbia **63**: 1-24.
- Brullo, S., Grillo, M. & Guglielmo, A. 1996: Considerazioni fitogeografiche sulla flora iblea. — Boll. Acc. Gioenia Sci. Nat. **29**: 45-111.
- Camarda, I. & Ballero, M. 1981: Studi sulla flora e la vegetazione di Capo Carbonara (Sardegna meridionale). I: La Flora. — Boll. Soc. Sarda Sci. Nat. **20**: 157-185.
- Carmignani, L., Coccozza, T., Ghezzi, C., Pertusati, P. C. & Ricci, A. C. 1982: Lineamenti del basamento sardo. Guida alla geologia del Paleozoico sardo. Mem. Soc. Geol. It. — Pp. 11-23 in: Guide Geologiche Regionali. — Cagliari.
- Chiappini, M. & Diana, G. 1978: Distribuzione geografica ed ecologica dell'endemismo sardo-corso *Brassica insularis* Moris in Sardegna. — Morisia **4**: 55-62.
- Corsi, G. 1963: Citologia, embriologia e distribuzione geografica di *Brassica insularis* Moris, endemismo sardo-corso. — Ann. Bot. (Roma) **27**: 419-430.
- Durán, M., Pinna, M. S., Orrù, M., Sanna, A., Fenu, G., Cogoni, D., Nebot, A. & Bacchetta, G. 2016: Conservación y restauración de los hábitats de dunas costeras en Cerdeña: la experiencia del proyecto LIFE+ PROVIDUNE. — In: Xavier Roig-Munar, F. (ed.) Restauración y gestión de sistemas dunares Estudios de casos. Recerca i Territori, **8**. — Torroella de Montgrí.

- Fenu, G., Fois, M., Cañadas, E. M. & Bacchetta, G. 2014: Using endemic-plant distribution, geology and geo-morphology in biogeography: the case of Sardinia (Mediterranean Basin). – *Syst. Biodivers.* **12**: 181-193.
- Galasso, G., Conti, F., Peruzzi, L., Ardenghi, N. M. G., Banfi, E., Celesti-Gradow, L., Albano, A., Alessandrini, A., Bacchetta, G., Ballelli, S., Bandini Mazzanti, M., Barberis, G., Bernardo, L., Blasi, C., Bouvet, D., Bovio, M., Cecchi, L., Del Guacchio, E., Domina, G., Fascetti, S., Gallo, L., Gubellini, L., Guiggi, A., Iamónico, D., Iberite, M., Jiménez-Mejías, P., Lattanzi, E., Marchetti, D., Martinetto, E., Masin, R. R., Medagli, P., Passalacqua, N. G., Peccenini, S., Pennesi, R., Pierini, B., Podda, L., Poldini, L., Prosser, F., Raimondo, F. M., Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A., Scortegagna, S., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagensommer, R. P., Wilhelm, T. & Bartolucci, F. 2018: An updated checklist of the vascular flora alien to Italy. – *Pl. Biosyst.* **152**: 556-592. <https://doi.org/10.1080/11263504.2018.1441197>
- Iiriti, G. 2006: Flora e paesaggio vegetale del Sarrabus-Gerrei (Sardegna sud orientale). – Tesi di Dottorato in Botanica Ambientale ed Applicata (XIX ciclo). – Cagliari.
- Ladero Alvarez, M., Díaz González, T. E., Penas Merino, A., Rivas-Martínez, S. & Valle Gutiérrez, C. 1987: Datos sobre la vegetación de las Cordilleras Central y Cantábrica. – *Itinera Geobot.* **1**: 3-147.
- Mayer, A. 1995: Comparative study of the coastal vegetation of Sardinia (Italy) and Crete (Greece) with respect to the effects of human influence. – *Libri Botanici* **15**. – München.
- Mayoral, O., Podda, L. & Porceddu, M. 2020: Invasive alien flora on coastal Mediterranean habitats. – In: Grigore, MN. (eds) *Handbook of Halophytes*. – Cham. https://doi.org/10.1007/978-3-030-17854-3_18-1.
- Moris, G. G. 1827: *Stirpium Sardoarum Elenchus*. – Carali.
- 1837-1859: *Flora Sardoia*, **1-3**. – Taurini.
- Mossa, L. & Fogu, M. C. 1987: La vegetazione dell'Isola dei Cavoli. – *Ann. Bot. (Roma)* **45**: 133-144.
- & Tamponi, G. 1978: La flora e la vegetazione dell'Isola dei Cavoli (Sardegna sud-orientale). – *Rend. Sem. Fac. Sci. Nat. Univ. Cagliari* **48**: 433-463.
- , Curreli, F. & Fogu, M. C. 2000: La vegetazione degli habitats terrestri della riserva marina protetta di Capo Carbonara (Sardegna sud-orientale). – *Rend. Sem. Fac. Sci. Univ. Cagliari* **70**: 163-185.
- , Guarino, R. & Fogu, M. C. 2003: Contributo alla conoscenza della flora terofitica della Sardegna. – *Rend. Sem. Fac. Sci. Univ. Cagliari* **73**: 1-209.
- Orrù, P., Cocco, A. & Panizza, V. 1994: Rilevamento geomorfologico subacqueo del settore compreso tra Capo Boi e Punta Is Cappuccinus (Sardegna Sud-Occidentale). – *Mem. Descr. Carta Geol. d'It.* **52**: 163-176.
- Palmese, M. T., Uncini Manganelli, R. & Tomei, P. E. 2001: An ethno-pharmacobotanical survey in the Sarrabus district (south east Sardinia). – *Fitoterapia* **72**: 619-643.
- Pignatti, S. 1982: *Flora d'Italia* **1-3**. – Edagricole, Bologna.
- Pinna, M. S., Cañadas, E. M., Fenu, G. & Bacchetta, G. 2015: The European *Juniperus* habitat in the Sardinian coastal dunes: Implication for conservation. – *Estuar. Coast. Shelf S.* **164**: 214-220.
- , Bacchetta, G., Orrù, M., Cogoni, D., Sanna, A. & Fenu, G. 2017: Results of the Providune project: restoration of the “Coastal dunes with *Juniperus* spp” priority habitat in Sardinia. – *Pl. Sociol.* **54**: 73-84.
- Podda, L., Lazzeri, V., Mascia, F., Mayoral García Berlanga, O. & Bacchetta, G. 2012: The Checklist of the Sardinian Exotic Flora: an Update. – *Not. Bot. Horti Agrob.* **40**: 14-21, + Annex I checklist, 1-11.

- , Santo, A., Mattana, M., Mayoral, O. & Bacchetta, G. 2018: Inter- and intra-variability of seed germination traits of *Carpobrotus edulis* N. E. Br. and its hybrid *C. affine acinaciformis*. – Pl. Biol. **20**: 1059-1067. <https://doi.org/10.1111/plb.12867>
- Puddu, S., Podda, L., Mayoral, O., Delage, A., Hugot, L., Petit, Y. & Bacchetta, G. 2016: Comparative analysis of the alien vascular flora of Sardinia and Corsica. – Not. Bot. Horti Agrob. **44**: 337-346.
- Pyšek, P., Richardson, D. M., Rejmánek, M., Webster, G. L., Williamson, M. & Kirschner, J. 2004: Alien plants in checklist and floras: towards better communication between taxonomist and ecologists. – Taxon **53**: 131-143.
- Raunkiaer, C. 1934: The life forms of plants and statistical plant geography. – Oxford.
- Richardson, D. M., Pyšek, P. & Carlton, J. T. 2011: A compendium of essential concepts and terminology in biological invasions. – Pp. 409-420 in: Richardson, D. M. (ed), Fifty years of invasion ecology: the legacy of Charles Elton. – Oxford.
- , —, Rejmánek, M., Barbour, M. G., Panetta, F. D. & West, C. J. 2000: Naturalization and invasion of alien plants: Concepts and definitions. – Divers. Distrib. **6**: 93-107.

Addresses of the authors:

Lina Podda¹, Francesca Meloni^{1*}, Giuseppe Fenu¹, Gianluca Iiriti² & Gianluigi Bacchetta^{1,2},

¹Centro Conservazione Biodiversità (CCB), Dipartimento di Scienze della Vita e dell'Ambiente (DISVA), Università degli Studi di Cagliari, V.le S. Ignazio da Laconi 13, 09123 Cagliari, Italy.

²Hortus Botanicus Karalitanus (HBK), Università degli Studi di Cagliari, V.le S. Ignazio da Laconi 9-11, 09123 Cagliari, Italy.

*Corresponding author. Email: meloni.francesca72@gmail.com

